

The Generalized Witt Algebras using Additive Maps

Ki-Bong Nam and Mook Ok Wang

February 8, 2008

Abstract

Kawamoto generalized the Witt algebra using $F[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ instead of $F[x_1, \dots, x_n]$. We construct the generalized Witt algebra $W(g_p, n)$ by using an additive map g_p from a set of integers into a field of characteristic zero where $1 \leq p \leq n$. We show that the Lie algebra $W(g_p, n)$ is simple if g_p is injective, and also the Lie algebra $W(g_p, n)$ has no ad-diagonalizable elements.

1 Introduction

Let F be a field of characteristic zero. The Witt algebra is called the general algebra by Rudakov [8]. Kac [2] studied the generalized Witt algebra on the F -algebra in the formal power series $F[[x_1, \dots, x_n]]$ for a positive integer n . Nam [5] constructs the Lie algebra on the F -subalgebra $F[e^{\pm x_1}, \dots, e^{\pm x_n}, x_1, \dots, x_{n+m}]$ in the formal power series $F[[x_1, \dots, x_{n+m}]]$ for the positive integers n and m .

The Witt algebra $W(n)$ has a basis

$$\{x_1^{a_1} \cdots x_n^{a_n} \partial_i | a_1, \dots, a_n \in \mathbb{N}, 1 \leq i \leq n\}$$

with Lie bracket on basis elements

$$\begin{aligned} & [x_1^{a_1} \cdots x_n^{a_n} \partial_i, x_1^{b_1} \cdots x_n^{b_n} \partial_j] \\ &= b_i x_1^{a_1+b_1} \cdots x_n^{a_n+b_n} x_i^{-1} \partial_j - a_j x_1^{a_1+b_1} \cdots x_n^{a_n+b_n} x_j^{-1} \partial_i \end{aligned}$$

where $\mathbb{N}$ is the set of non-negative integers.

Consider the generalized Witt algebra $W(n, m)$ having a basis

$$\{e^{a_1 x_1} \cdots e^{a_n x_n} x_1^{u_1} \cdots x_{n+m}^{u_{n+m}} \partial_k | a_1, \dots, a_n, u_1, \dots, u_{n+m} \in \mathbb{Z}, 1 \leq k \leq n+m\}$$

with Lie bracket on basis elements given by

$$\begin{aligned} & [e^{a_1 x_1} \cdots e^{a_n x_n} x_1^{u_1} \cdots x_{n+m}^{u_{n+m}} \partial_i, e^{b_1 x_1} \cdots e^{b_n x_n} x_1^{t_1} \cdots x_{n+m}^{t_{n+m}} \partial_j] \\ &= b_i e^{a_1 x_1 + b_1 x_1} \cdots e^{a_n x_n + b_n x_n} x_1^{u_1+t_1} \cdots x_{n+m}^{u_{n+m}+t_{n+m}} \partial_j \end{aligned}$$

$$\begin{aligned}
& +t_i e^{a_1 x_1 + b_1 x_1} \dots e^{a_n x_n + b_n x_n} x_1^{u_1 + t_1} \dots x_{n+m}^{u_{n+m} + t_{n+m}} x_i^{-1} \partial_j \\
& -a_i e^{a_1 x_1 + b_1 x_1} \dots e^{a_n x_n + b_n x_n} x_1^{u_1 + t_1} \dots x_{n+m}^{u_{n+m} + t_{n+m}} \partial_i \\
& -u_i e^{a_1 x_1 + b_1 x_1} \dots e^{a_n x_n + b_n x_n} x_1^{u_1 + t_1} \dots x_{n+m}^{u_{n+m} + t_{n+m}} x_j^{-1} \partial_i
\end{aligned}$$

where $b_i = 0$ if $n + 1 \leq i \leq n + m$, and $a_j = 0$ if $n + 1 \leq j \leq n + m$ (see [4, 5, 7]). In [5, 8], it is noted that the Lie subalgebra of $W(n, m)$ is the Witt algebra $W(n)$ on $F[x_1, \dots, x_m]$. Let g_p be an additive map from Z into F where $1 \leq p \leq n$. Let us define as $W(g_p, n)$ the Lie algebra with basis

$$B := \left\{ \binom{a_1}{i_1} \dots \binom{a_n}{i_n}_k \mid a_1, \dots, a_n, i_1, \dots, i_n \in Z, 1 \leq k \leq n \right\} \quad (1)$$

and a Lie bracket on basis elements given by

$$\begin{aligned}
& \left[\binom{a_1}{i_1} \dots \binom{a_n}{i_n}_k, \binom{b_1}{j_1} \dots \binom{b_n}{j_n}_l \right] \\
& = g_k(b_k) \binom{a_1 + b_1}{i_1 + j_1} \dots \binom{a_n + b_n}{i_n + j_n}_l \\
& + j_k \binom{a_1 + b_1}{i_1 + j_1} \dots \binom{a_k + b_k}{i_k + j_k - 1} \binom{a_{k+1} + b_{k+1}}{i_{k+1} + j_{k+1}} \dots \binom{a_n + b_n}{i_n + j_n}_l \\
& - g_l(a_l) \binom{a_1 + b_1}{i_1 + j_1} \dots \binom{a_n + b_n}{i_n + j_n}_l \\
& - i_l \binom{a_1 + b_1}{i_1 + j_1} \dots \binom{a_l + b_l}{i_l + j_l - 1} \binom{a_{l+1} + b_{l+1}}{i_{l+1} + j_{l+1}} \dots \binom{a_n + b_n}{i_n + j_n}_k.
\end{aligned} \quad (2)$$

It follows from [1, 4, 5, 6] that the above bracket is extended linearly to the given basis B . Also, it is not hard to show that the above bracket satisfies the Jacobi identity. In section 2, we will prove the following main theorems. Throughout the paper, a given map g_p means an additive and injective map from a set of integers into a field of characteristic zero.

Theorem 1 The Lie algebra $W(g_p, n)$ is simple.

Theorem 2 For any automorphism of $\theta \in \text{Aut}(W(g, 1))$

$$\theta \left(\binom{0}{1}_1 \right) = \sum_j C_j \left(\binom{0}{j}_1 \right)$$

where $C_j \in F$.

Theorem 3 Each derivation of $W(g, 1)_+$ can be written as a sum of an inner derivation and a scalar derivation [6].

2 Simplicities of $W(g_p, n)$

The Lie algebra $W(g_p, n)$ has a Z^n -gradation as mentioned in [3] : that is,

$$W(g_p, n) = \bigoplus_{(a_1, \dots, a_n) \in Z^n} W_{(a_1, \dots, a_n)} \quad (3)$$

where each is the subspace of $W(g_p, n)$ with a basis

$$\bar{B} = \left\{ \binom{a_1}{i_1} \cdots \binom{a_n}{i_n} \mid a_1, \dots, a_n, i_1, \dots, i_n \in Z, 1 \leq l \leq n \right\}$$

Let $W_{(a_1, \dots, a_n)}$ denote the $(a_1, \dots, a_n)$ -homogeneous component of $W(g_p, n)$ and elements in the $W_{(a_1, \dots, a_n)}$ -homogeneous $(a_1, \dots, a_n)$ -homogeneous elements. Note that the $(0, \dots, 0)$ - homogeneous component is isomorphic to the Witt algebra $W(n)$ [8]. From now on let $(0, \dots, 0)$ -homogeneous component denote the 0-homogeneous component. For the simplicity of $W(g_p, n)$, we define the map as an additive and injective map. Now we introduce a lexicographic ordering of two basis elements of $W(g_p, n)$ as follows : for any two elements $\binom{a_1}{i_1} \cdots \binom{a_n}{i_n}_l, \binom{b_1}{j_1} \cdots \binom{b_n}{j_n}_k$, we have

$$\binom{a_1}{i_1} \cdots \binom{a_n}{i_n}_l > \binom{b_1}{j_1} \cdots \binom{b_n}{j_n}_k$$

if $(a_1, \dots, a_n, i_1, \dots, i_n, l) > (b_1, \dots, b_n, j_1, \dots, j_n, k)$ by the natural lexicographic ordering in Z^{2n+1} .

For any element $l \in W(g_p, n)$, l can be written as follows using the ordering and gradation :

$$\begin{aligned} l = & \sum_{i_1, \dots, i_n, p} C(i_1, \dots, i_n, p) \binom{a_{11}}{i_1} \cdots \binom{a_{1n}}{i_n}_l + \cdots \\ & + \sum_{j_1, \dots, j_n, q} C(j_1, \dots, j_n, q) \binom{a_{t1}}{j_1} \cdots \binom{a_{tn}}{j_n}_q \end{aligned}$$

where $1 \leq p, \dots, q \leq n$, $(a_{11}, \dots, a_{1n}) > \dots > (a_{t1}, \dots, a_{tn})$ and

$$C(i_1, \dots, i_n, p), \dots, C(j_1, \dots, j_n, q) \in F.$$

Next, define the string number $st(l) = t$ for l (see [5, 6]), and $l_p(l)$ as $\max\{i_1, \dots, i_n, \dots, j_1, \dots, j_n\}$. For any basis element $\binom{a_1}{i_1} \cdots \binom{a_n}{i_n}_l$ in $\bar{B}$, let us refer to $a_1, \dots, a_n$ as upper indices and $i_1, \dots, i_n$ as lower indices.

Remark 2.1 If g_p is an inclusion, then $W(g_p, n) = W(n, 0)$ is the generalized Witt algebra which is studied by Nam [5, 6].

Lemma 1 *If l is any non-zero element, then the ideal $\langle l \rangle$ generated by l contains an element whose lower indices are positive.*

Proof. Let l be a nonzero element of $W(g_p, n)$. Take an element $M = \binom{0}{j_1} \cdots \binom{0}{j_n}_t$ such that $j_1 \gg \cdots \gg j_n$ and t such that either $a_t \neq 0$ or $i_t \neq 0$ in l , where $a \gg b$ means a is sufficiently larger than b . Then $0 \neq [M, l]$ is the required element. $\square$

Lemma 2 *If an ideal I of $W(g_p, n)$ contains $\binom{0}{0} \cdots \binom{0}{0}_l$ for $1 \leq l \leq n$, then $I = W(g_p, n)$.*

Proof. Since $W(n)$ is a simple subalgebra of $W(g_p, n)$ with the basis $\{\binom{0}{i_1} \cdots \binom{0}{i_n}_l | i_1, \dots, i_n \in Z, 1 \leq l \leq n\}$, the ideal $\langle \binom{0}{0} \cdots \binom{0}{0}_i \rangle$ contains $W(n)$, where $\langle \binom{0}{0} \cdots \binom{0}{0} \rangle$ is the ideal of which is generated by $\binom{0}{0} \cdots \binom{0}{0}_l$ for a fixed $t \in \{1, \dots, n\}$ (see [8]). It follows from Lemma 2 of [5] and Lemma 3 of [6] that the basis elements of $W(g_p, n)$ are contained in I by using the injectiveness of g_p . $\square$

Theorem 1 *The Lie algebra $W(g_p, n)$ is simple.*

Proof. It is not difficult to prove this theorem by induction on $st(l)$ for any element l in any ideal I of $W(g_p, n)$ and by considering the one to one properties of g_p .

Corollary 1 *The Lie algebra $W(n, 0)$ is simple.*

Proof. If we take an additive embedding $g_p : Z \rightarrow F$, $1 \leq p \leq n$, then we get the required result (see [5, 6]). $\square$

It is an interesting problem to find all the automorphisms of the subalgebra $W(g, 1)$ of $W(g_p, n)$.

Theorem 2 *For any automorphism $\theta \in \text{Aut}(g_p, n)$*

$$\theta\left(\binom{0}{1}_1\right) = \sum_j C_j \binom{0}{j}_1$$

where $C_j \in F$.

Proof. It is not difficult to prove this theorem using the gradation of (2.1) and the $W(g, 1)$ acting of $\binom{0}{0}_1$ on the zero homogeneous component W_0 whose vector basis is $\{\binom{0}{i}_1 | i \in Z\}$ as an ad-map. $\square$

If we consider the Lie subalgebra of $W(g, 1)$ such that all the lower indices are zero, then this subalgebra is a Block algebra $W(1)$ which is also called the centerless Virasoro algebra [1]. Thus all the automorphisms of this Lie algebra can be determined by Theorem 3 in [1].

Consider the Lie subalgebra $W(1, 1)^+$ of $W(n, m)$ in [6] with basis

$$\{e^{ax} x^i y^j \partial_x, e^{bx} x^l y^m \partial_y | a, b \in Z, i, j, l, m \in N\}$$

where N is the set of non-negative integers.

Conjecture For any $\theta \in \text{Aut}(W(1, 1)^+)$ $\theta(y\partial_y) = (\alpha y + \beta)\partial_y$ for some $0 \neq \alpha, \beta \in F$.

The element $l \in W(g_p, n)$ is an ad-diagonalizable element if $[l, m] = \alpha(m)m$ for any $m \in B$ given in (1) and for some $\alpha(m) \in F$.

Proposition 1 *The Lie algebra $W(g, n)$ has no ad-diagonalizable element with respect to the basis given in (1).*

Proof. Since $W(g, n)$ is Z^n -graded Lie algebra all the ad-diagonalizable elements are in the $(0, \dots, 0)$ -homogeneous component. $W_{(0, \dots, 0)}$ is isomorphic to $W(n)$ as Lie algebras where $W(n)$ is the well known Witt algebra [8]. Thus all the ad-semisimple elements of $W(g_p, n)$ are of the form $\sum_{i=1}^n C_i \binom{0}{1}_i$, where $C_i \in F$. But $[\sum_{i=1}^n C_i \binom{0}{1}_i, \binom{a}{0}_j] \neq \alpha \binom{a}{0}_j$ for any $\alpha \in F$. Therefore, we proved the proposition. $\square$

Remark 2.2 For another proof of Proposition 2.6, see Corollary 1 of [5].

3 Derivation of $W(g, 1)_+$

Consider the subalgebra $W(g, 1)_+$ of $W(g_p, n)$ with basis

$$B_+ := \left\{ \binom{a}{i}_1 \mid a \in Z, i \in N \right\}$$

where N is the set of non-negative integers.

In this section we determine all the derivations of the Lie algebra $W(g, 1)_+$. Ikeda and Kawamoto found all the derivations of the Kawamoto algebra $W(G, I)$ [4] in their paper [2]. It is very important to find all the derivations of a given Lie algebra to compare with other Lie algebras. Let L be a Lie algebra over any field F . An F -linear map D from L to L is a derivation if $D([l_1, l_2]) = [D(l_1), l_2] + [l_1, D(l_2)]$ for any $l_1, l_2 \in L$.

Let L be a Lie algebra over any field F . Define the derivation D of L to be a scalar derivation if for all basis elements l of L , $D(l) = f_l l$ for some scalar $f_l \in F$ [1], [7].

We need the following lemma.

Lemma 3 *Let D be a derivation of $W(g, 1)_+$. If $D(\partial) = 0$, then $D = f \text{ad}_\partial + S$ where $f \in F$ and S is a scalar derivation.*

Proof. It is not difficult to prove this lemma using the gradation and the ordering of the Lie algebra $W(g, 1)_+$. [6]. $\square$

Theorem 3 *Each derivation of $W(g, 1)_+$ can be written as a sum of an inner derivation and a scalar derivation.*

Proof. Let D be any derivation of $W(g, 1)_+$. Then $D(\partial) = f\partial$ for some $f \in F[e^{\pm x}, x]$. Since $\partial : F[e^{\pm x}, x] \rightarrow F[e^{\pm x}, x]$ is onto, there is a function $g \in F[e^{\pm x}, x]$ such that $\partial(g) = f$. Then $ad_{g\partial}(\partial) = [\partial, g\partial] = f\partial = D(\partial)$. We have $(D - ad_{g\partial})(\partial) = 0$. By Lemma 3 we have $D = ad_{g\partial} + cad_{\partial} + S$ [6]. Therefore, we have proved the theorem. $\square$

Remark 3.1 Let Z^+ and F^+ be the additive group of Z and F respectively. For any $g \in Hom(Z^+, F^+)$, we have $g(1) = m$ for some fixed $m \in F$, thus the Witt algebra $W(n)$ can not be changed by using the idea of the additive map in this paper [5].

References

- [1] R. Block, *On torsion-free abelian groups and Lie algebras*, Proc. Amer. Math. Soc. 9(1958), 613-620.
- [2] T. Ikeda and N. Kawamoto, *On the derivations of generalized Witt algebras*, Hiroshima Math. J., 20 (1990), 47-55.
- [3] V. G. Kac, *Description of filtered Lie algebra with which graded Lie algebras of Cartan type are associated*, Izv. Akad. Nauk SSSR, Ser. Mat. Tom. 38(1974), 832-834.
- [4] N. Kawamoto, *On G-Graded Automorphisms of generalized Witt algebras*, Contem. Math. A.M.S., 184(1995), 225-230.
- [5] N. Kawamoto, *Generalized Witt algebras over a field of characteristic zero*, Hiroshima. Math. J., 16(1986), 417-426.
- [6] Ki-Bong Nam, *Generalized Witt algebras over a field of characteristic zero*, UW-Madison, thesis, (1998), 1-40.
- [7] Ki-Bong Nam, *Simple Lie algebras which generalize Witt algebras I*, Kyungpook Math. J., to appear, 1-9.
- [8] D. S. Passman, *Simple Lie Algebras of Witt-Type*, to appear, 1-9.
- [9] A. N. Rudakov, *Groups of Automorphisms of Infinite-Dimensional Simple Lie Algebras*, Math. USSR-Izv. 3(1969), 707-722.

Department of Mathematics, UW-Madison, WI 53706

e-mail :nam@math.wisc.edu

Department of Mathematics, Hanyang University-Ansan, Ansan, Korea